

UNSTABLE PARTICLES AND GAUGE INVARIANCE

Geert Jan van Oldenborgh

Paul Scherrer Institute, CH-5232 Villigen PSI, Switzerland

February 7, 2008

Abstract

When computing the properties of reactions involving unstable charged particles care has to be taken to use a gauge invariant amplitude. In this talk we present methods to (automatically) obtain such an amplitude, both at the tree level and in one-loop calculations, using only a minimal number of diagrams. The numerical difference with gauge variant methods commonly used will be discussed for the case of tree level W pair production.

PSI-PR-93-19

1 Introduction

Many of the cross sections one would like to evaluate with the methods described elsewhere in these proceedings involve the production and decay of unstable particles. The W and Z bosons both have a sizable width Γ ($\Gamma/m \approx 1/40$), and the same may be true for the top quark and Higgs boson (depending on their masses). For these particles the well-known narrow width approximation will not be sufficient for many purposes. Including off-shell effects, however, involves a resummation of some of the higher order graphs, which means that care has to be taken to keep the matrix element gauge invariant. This last point is not only of academic interest as gauge breaking terms are often much larger numerically than the gauge invariant result. Therefore even suppression factors Γ/m may not be enough to avoid inaccuracies.

The Dyson resummation of higher order graphs in the propagator, which leads to a finite width, is clearly only needed when the unstable particle can kinematically be on its mass shell. Indeed, for an unstable particle with space-like momentum $p^2 < 0$ the imaginary part of the self energy is zero, hence no finite width should be used. One can therefore not use a complex mass in the Lagrangian, even disregarding the problems with the standard model mass generation mechanism which would follow (for instance a complex value for $\sin^2 \theta_W$). It is a purely kinematical problem which particles should be given a finite width and which not.

The amplitude can thus be divided into a set of *resonant* diagrams, which contain a divergence within the allowed phase space as an unstable particle is taken on-shell, and *non-resonant* graphs without such divergences. In principle the contribution of the non-resonant diagrams is suppressed with respect to the resonant ones by $\Gamma/m \propto \alpha$. We first discuss ways to only use the resonant tree graphs to get the dominant behaviour, next increasing the accuracy of the result by including the non-resonant tree graphs and the resonant one-loop graphs. A more detailed discussion will be given in a forthcoming publication [1]. The tree level results are based on Ref. [2].

2 Resonant Tree Graphs

In this section some schemes will be given to deal with the resonant tree graphs. For simplicity these are first discussed for a single unstable particle with momentum p and mass m ; the extension to more particles will be given later later. We assume that the diagram generator has already sorted out which diagrams ones are resonant and which are not. Without resummation the full (gauge invariant) amplitude for this process therefore has the form

$$\mathcal{M} = \frac{R(p^2)}{p^2 - m^2} + N(p^2) . \quad (1)$$

The essential variable here is the virtuality p^2 of the unstable particle. The other kinematical variables are assumed to be CMS angles, which are independent of p^2 . Unfortunately, this amplitude gives an infinite cross section when the integration region includes the pole at $p^2 = m^2$.

The simplest approximation one can make is to completely factorize the process in production and decay: the narrow width approximation. This fails to take into account terms of $\mathcal{O}(\Gamma)$; the corresponding dimensionless parameter will usually be Γ/m (about $1/40$ for the W), but can be much larger near the threshold for production of the unstable particles.

The most important of these $\mathcal{O}(\Gamma)$ corrections are included by simply using the on-shell expression for the matrix element, but treating the kinematics and unstable particle propagator off-shell. This off-shell propagator is derived by resumming the one-loop corrections to the propagator, giving rise to a simple Breit-Wigner propagator $1/(p^2 - m^2 + im\Gamma)$ with Γ defined by the relation $m\Gamma = \text{Im}\Pi(m^2)$ in terms of the self energy Π evaluated at the real mass. We refer to this procedure as the “resonant” scheme. This procedure obviously respects gauge invariance and will serve as the base line to compare the others with.

The next step is usually to also evaluate the matrix element off-shell, we refer to this procedure as the “off-shell” method. This also gives an answer below threshold. In the case of a charged resonance however, this procedure violates $U(1)_{em}$ gauge invariance, even if the width is given by the physical (on-shell) quantity, which is gauge invariant. The reason is that the original resonant graphs are not the only graphs which lead to the particular final state: non-resonant graphs have to be included.

3 Non-resonant Tree Graphs

One reason to include the non-resonant graphs is to conserve gauge invariance. Another is that they can be sizable due to some other near-divergence, such as large collinear logarithms. The “naïve” way to include these non-resonant graphs is to just add them to the resonant terms with a complex mass. However, as the amplitude was gauge invariant for $\Gamma = 0$ it follows that it must be gauge variant for $\Gamma > 0$ (if the resonant graphs are not separately gauge invariant).

An obviously gauge invariant procedure to include the non-resonant graphs was introduced by Zeppenfeld *et al* [3]. The price to save gauge invariance in this “overall” scheme is an incorrect treatment of the non-resonant contribution close to mass shell. However, it can be argued that the difference is of higher order in this region of phase space. We thus have a prescription which is correct to leading order both on resonance and away from it, but these contributions are formally of different order in α .

Another gauge invariant way to include the off-shell effects is to systematically separate orders in Γ [4]. As the residue at the pole $p^2 = m^2$ is gauge invariant one can add the finite width in the first term without breaking gauge invariance. This corresponds to adding and resumming only the gauge invariant part of the propagator corrections. In this scheme the cross section is given as the sum of the

name	kine- matics	matrix element	gauge invari- ant?	treatment non-resonant terms	threshold behaviour
narrow width	on-shell	$R(m^2)$	yes	none	undefined be- low threshold
resonant	off-shell	$\frac{R(m^2)}{p^2 - m^2 + im\Gamma}$	yes	none	zero below threshold
off-shell	off-shell	$\frac{R(p^2)}{p^2 - m^2 + im\Gamma}$	no	none	wrong below threshold
naive	off-shell	$\frac{R(p^2)}{p^2 - m^2 + im\Gamma} + N(p^2)$	no	ok	ok
overall	off-shell	$\frac{R(p^2)}{p^2 - m^2 + im\Gamma} + \frac{p^2 - m^2}{p^2 - m^2 + im\Gamma} N(p^2)$	yes	wrong	ok
pole- scheme	off-shell	$\frac{R(m^2)}{p^2 - m^2 + im\Gamma} + \frac{R(p^2) - R(m^2)}{p^2 - m^2} + N(p^2)$	yes	ok	wrong around threshold

Table 1: The six different schemes to treat unstable particles at tree level.

resonant cross section plus $\mathcal{O}(\Gamma)$ corrections, which are both gauge invariant. It is thus a natural starting point for higher order corrections.

However, this “polescheme” also has some undesirable properties. The first (resonant) term has a discontinuity when the threshold for the production of the unstable particle is crossed. Approaching from below, one even encounters the original (non-resummed) singularity in the propagator. The accuracy of this scheme is thus doubtful around threshold.

The difference between the last three schemes is of order Γ^2 for total cross sections and singly differential distributions, but of order Γ for doubly and higher differential distributions. This follows from the occurrence of a Levi-Civita tensor in the $\mathcal{O}(\Gamma)$ terms. A summary of all schemes is given in table 1.

The extensions to the case where there are more unstable particles are largely straightforward. There now appears a hierarchy of n -fold resonant diagrams. The first three methods just use the maximally resonant ones. In the “polescheme” one only needs these and the ones where only one particle is non-resonant; the “overall” scheme uses all diagrams. The threshold behaviour of the ‘polescheme’ now is worse than before. However, in this area the worth of any fixed-order calculation is doubtful because of possible bound state effects.

The numerical influence on the total cross section is illustrated in Fig. 1, where the total cross section for $\gamma\gamma \rightarrow \ell^- \bar{\nu}_\ell W^+$ is plotted, with the W^+ considered stable and a photon beam derived from backscattered electrons [5]. The physically relevant off-shell effects will be twice as large. To avoid the collinear divergence in the final state a minimum transverse momentum $p_\perp(\ell) > 0.02\sqrt{s}$ is demanded; this should also give an indication of detector acceptance. It can be seen that the narrow width approximation misses some important effects. The graph for the offshell scheme is misleading, in that we used the formula $\sum_i \epsilon_i^\mu \epsilon_i^\nu = -g^{\mu\nu}$ which is not valid when the photon couples to a non-conserved current. It does show that one has to be careful when breaking gauge invariance. One should also note that the differences between the various schemes are expected to be much larger in doubly differential distributions.




Figure 1: Cross sections for $\gamma\gamma \rightarrow \ell^- \bar{\nu}_\ell W^+$ in the different schemes defined in the text with $p_\perp(\ell) > 0.02\sqrt{s_{ee}}$

4 Resonant One-loop Corrections

When computing the $\mathcal{O}(\alpha)$ corrections one is normally only interested in the resonant part. We will thus have to identify which diagrams contribute to this, and then to find a gauge invariant way to isolate just the resonant part. A generalization of the “polescheme” is proposed for this last step. In the case of a neutral resonance (like the Z boson) the isolation step is not necessary, as the resonant diagrams already form a gauge invariant subset; the only bone of contention left in this case amounts to the proper definition of the mass of the unstable particle.

There are three classes of resonant diagrams. The first one contains the corrections to the resonant propagator. A gauge invariant part of these are resummed to give the finite width; the rest (vanishing at $p^2 = m^2$) has to be treated perturbatively. The second group, the factorizable diagrams, comprises the corrections to either production or decay of an unstable particle; these diagrams obviously retain the original resonant propagators. Added to this, charged resonances will give rise to logarithms $\log(p^2 - m^2)$ as the virtuality and finite width of the unstable particle regulate the infrared divergence. The third class of non-factorizable diagrams is given by the set of infrared divergent diagrams with the photon (or gluon) spanning one or more resonant propagators. Simple power counting arguments show that no other non-factorizable diagrams can give a resonant contribution. One would suspect that even the contributions from these diagrams may cancel against the corresponding soft Bremsstrahlung diagrams. As the hard Bremsstrahlung is easily seen to be non-resonant this would imply the intuitively satisfying result that initial-final and final-final state interaction vanish in the narrow width approximation.

However, this is not the case. As an example consider the doubly resonant five point function shown in Fig. 2. Using the Schouten identity one can decompose it




Figure 2: A double resonant five point function occurring in W pair production

in five four point functions as

$$E_0 = \frac{-1}{2\Delta_{5s}} \left(\Delta_{ps}^\gamma D_0(\gamma) + \Delta_{ps}^\mu D_0(\mu) + \Delta_{ps}^{W^-} D_0(W^-) + \Delta_{ps}^{W^+} D_0(W^+) + \Delta_{ps}^e D_0(e) \right) \quad (2)$$

where $D_0(p)$ denotes the scalar four point function with the propagator corresponding to particle p missing and the Δ 's denote determinants of external and internal momenta [6]. The overall factor $-1/2\Delta_{5s}$ is quadratic in the virtualities $p_\pm^2 - m^2$ of the W bosons and thus gives rise to the double resonance. The $D_0(\gamma)$ and Δ_{ps}^γ are both finite, whereas for the other combinations the determinant is linear in the virtuality and the four point function linearly divergent.

The corresponding scalar Bremsstrahlung integral can be decomposed similarly, except that the equivalent of the $D_0(\gamma)$ never appears. The other four terms exactly match the corresponding terms in the virtual diagram, leaving this one doubly resonant term. There is no symmetry which would cause it to disappear, even when integrated over phase space, and there are no other graphs which will give rise to the same four point function. It therefore is a double resonant term of order α which is not suppressed by additional factors Γ/m . The failing cancellations can be traced back to the fact that, in the limit $\Gamma \ll \lambda$ (λ is an infrared regulator, normally the photon mass) the linear and quadratic infrared divergences do *not* cancel. Only a numerical study can decide whether the non-factorizable diagrams give rise to measurable effects.

The diagrams we have selected as resonant diagrams in general do not form a gauge invariant subset. One can select a gauge invariant part by taking the residue at the pole, as was done in the previous sections. Unfortunately, this residue is ill-defined due to the occurrence of the divergent logarithms. (These also do not cancel between the virtual and Bremsstrahlung graphs.) These logarithms only occur in the scalar functions however. As the coefficients of the scalar functions are gauge invariant, one can evaluate these any way one likes.

The following recipe therefore serves as the extension of the “polescheme” for one-loop (and soft Bremsstrahlung) diagrams:

- The reduction of the diagram to scalar functions should be done on-shell (if the unstable particle is charged).
- The scalar functions can be evaluated in any convenient non-divergent way; on-shell or off-shell. This also allows one to take into account the finite width in non-divergent but non-analytical logarithms like the Coulomb singularity and threshold effects.

This schemes have been implemented in the FF library [7].

5 Conclusion

In this talk we have presented a scheme to systematically include off-shell effects in amplitudes including unstable particles, explicitly keeping gauge invariance. Unfortunately it breaks down near the threshold for the production of the unstable particles. In this region, however, the occurrence of bound state effects makes any

fixed-order perturbation series suspect. The numerical difference of this procedure with the (gauge variant) ones commonly used are small in the tree level total cross section, at least in the case of W pair production and the unitary gauge. Larger differences are expected for doubly differential cross sections.

6 Acknowledgements

The results presented here are based on work done together with Daniel Wyler and Andre Aeppli in the context of the one-loop corrections to W pair production.

References

- [1] A. Aeppli, G. J. van Oldenborgh and D. Wyler. (to appear).
- [2] A. Aeppli, F. Cuyppers and G. J. van Oldenborgh. Phys. Lett. **B314** (1993) 413.
- [3] D. Zeppenfeld, J. A. M. Vermaseren and U. Baur. Nucl. Phys. **B375** (1992) 3.
- [4] R. G. Stuart. Phys. Lett. **B262** (1991) 113, Phys. Lett. **B272** (1991) 353.
- [5] I. F. Ginzburg, Kotkin G. L., V. G. Serbo and V. I. Telnov. Nucl. Instr. Meth. **205** (1983) 47.
- [6] G. J. van Oldenborgh. One-loop Calculations with Massive Particles. Ph.D. thesis, Universiteit van Amsterdam, 1991.
- [7] G. J. van Oldenborgh. Comp. Phys. Comm. **66** (1991) 1.